

## FACTORS ASSOCIATED WITH THE INCIDENCE OF NEONATAL ASPHYXIA IN NEWBORNS AT THE GUIDO VALADARES NATIONAL HOSPITAL DILI TIMOR-LESTE

Ermelinda da Costa Monteiro\*, Carlos Boavida Tilman, José Dionísio Ximenes, Eugenia Carvalho de Araújo, Elisa de Deus, Etelvina José Tilman.

### *Abstract*

**Introduction:** Perinatal asphyxia is considered one of the major causes of neonatal deaths and neurological sequelae in newborns, resulting from maternal-fetal conditions or complications during childbirth. The report by the Guido Valadares de Dili National Hospital showed that 201 newborns died after birth and that 151 died at the age of 0-6 days and 8 died aged 7 to 28 days.

**Objective:** To analyze the factors associated with the incidence of neonatal asphyxia at the Guido Valadares National Hospital.

**Method:** Reconducted a descriptive and analytical study of quantitative approach, collecting a sample of 100 records of newborns of mothers with risk factors and used the Grid of Observation on Register of the midwife and the doctor.

**Results:** As maternal risk factors, 61% of mothers have parity higher than four children, and maternal pathologies in the sample include Hypertension 37%, Preeclampsia/Eclampsia 25%, Anemia 25% and Diabetes 13%. Regarding Hypertension, with mild degree 63%, and severe 38%; Preeclampsia/severe eclampsia 60% and mild 40%; in the degree of Mild Anemia (7 to 9 g/dl) 88%, and severe (<7g/dl) 12%; Type I Diabetes is 54% and Gestational Diabetes 46%; finally, premature membrane rupture is found in 51% of the sample. As for fetal risk factors for neonatal asphyxia, polyhydrosis 30%, Man Condon 13%, Infection 31%, Weight <1500 grams 13%, Inadequate weight for gestational age 28%. In the risk factors associated with delivery, we verified Forceps Delivery 13%, Prolapse of the cord 9% and Short Umbilical Cord 8%. In the evaluation of APGAR indices at the 1 minute verified or severe in 48 newborns 62% and mild 39%; at 5<sup>a</sup> minute, the APGAR Index was grave at 59% and 41%, lastly, it was found that in the APGAR index at 10 minutes, then Gave classification decreased 4%, and leve or moderate 7%. At 10 minutes 89% of the sample.

**Conclusion:** Hospitals, health centers and SISCa should promote the four minimum visits for pregnant women, in order to create a program of activities, such as home visits to define the rich factors of mothers. Pregnant women need to deliver with midwives and mothers in a controlled and safe environment to reduce the neonatal and post-neonatal mortality rate.

**Keyword:** Neonatal asphyxia, Maternal factors, fetal factors, factors of the work of the pair to and neonatal fafores, National Hospital Guido Valadares.

### INTRODUCTION

From a practical point of view, perinatal asphyxia can be understood as a failure in the establishment of normal breathing during the period of

birth, due to the impediment of oxygenation, at the time of delivery and in the period of expulsion. Currently, maternal and infant mortality, as an indicator of public health, is an important

global issue. One of the health problems that occur frequently at the time of birth of the baby is infant mortality as a result of asphyxiation<sup>1</sup>. If immediate nursing and midwives care is not provided to newborns with neonatal asphyxia, they will have a low probability of survival. When the newborn has neonatal asphyxia, the baby cannot breathe spontaneously and regularly, and it is found that babies who experience fetal distress before birth usually suffer from asphyxiation<sup>2</sup>. The main causes of neonatal mortality are intrinsically related to health and care received before, during and after childbirth. Neonatal asphyxia and birth trauma are usually caused by lack of medical follow-up and lack of access to obstetric specialists, such as midwives who help at home cited by Carlos & Ermelinda, 2022.

According to a world health organization (WHO) vision, every year 120 million babies born worldwide, out of a total of 4 million (33 per 1,000), die at the age of 30 days (neonatal) and about 3.6 million (3%) of these babies have neonatal asphyxia newborn babies in all world including Timor-Leste<sup>4</sup>. Mozambique, São Tomé and Príncipe, Timor-Leste, Guinea-Bissau and Brazil were evidenced by the decline in their infant mortality rates<sup>5</sup>. This is justified, perhaps, by the existence in our Constitution of the Democratic Republic of Timor-Leste in 2002, Part II, Title I, Article 18, on the protection of the child. The child is entitled to special protection by the family, the community and the Timor-Leste State cited by Carlos & Er-

melinda, 2022<sup>6</sup>. The 2009-2010 Public Health Survey D in Timor-Leste shows that the maternal mortality rate is 557 per 100,000 live births<sup>7</sup>. Based on this report of Statistics in Health carried out by the Office of Information System in Health and Epidemiologic Surveillance under Ministry of Health, in the year 2018, the fertility rate is 5.7 children per woman of reproductive age, the rate infant mortality is 44 per 1,000 live births and the post-neonatal and neonatal mortality rate is 22 per 1,000 live births. In the period of 2015, the number of neonatal mortalities, more specifically related to the age of 0 to 7 days, in health centers and in the reference and national hospitals currently existing in Timor-Leste, is of 119. In more detail you can see the data of each municipality: the Municipality of Dili with 36 dead neonates, Viqueque with 23, Bobonaro with 18, Ermera with 14, Lautem 13, Liquiça 12 and Ainaro 11. If we make a comparison between the years 2015 and 2016, in relation to the number of neonates killed in the municipality of Dili, we find that there was an increase of 73, in the number of deaths<sup>8</sup>.

In the same document, the reports of the hospitals and health centers for the year 2015, report that 999 newborn children had a weight of less than 2500 grams, and 203 died after birth. Of the latter, 154 died aged 0-6 days and 9 died aged 7-28 days, that is, in the neonatal period<sup>8</sup>. The report of the National Hospital Guido Valadares de Dili, showed that 925 babies were born weighing

less than 2500 grams, of which 127 died after birth. Of these, 100 died at the ages of 0-6 days.<sup>9</sup> Based on the data presented above, we verified a high infant mortality rate in the Hospital, so we propose to research the factors associated with the intention of neonatal asphyxia in the Guido Valadares National Hospital of Dili Timor-Leste.

### Goal

General objective: To analyze the factors associated with the incidence of neonatal asphyxia or suffocation at the Guido Valadares Dili Timor-Leste National Hospital.

Specific objectives are:

- Identify the incidence of neonatal asphyxia or suffocation;
- Describe the factors associated with neonatal suffocation or asphyxia related to the woman, mother and newborn fetus or baby;
- To verify the relationship of neonatal suffocation or asphyxia with the work performed by the midwife who attends the deliveries or newborns.

### THEORETICAL FRAMEWORK


Asphyxia is a baby state in which it does not breathe spontaneously and regularly at birth<sup>10</sup>. However, babies often experience asphyxiation only after childbirth. This problem is probably related to maternal health status, umbilical cord, problems with childbirth of problems in infants. Asphyxia means, etymologically, short-

ness of breath the u shortness of breath. Clinically it is a syndrome characterized by spon- sion the severe decrease of gas exchange oxo at the level of placenta of dealings, which results in hypoxemia, hypercapnia and tissue hypoxia, with metabolic acidosis<sup>11</sup>. Perinatal asphyxia is a problem suffered by the fetus or newborn (NB), due to monooxygenation or mal-perfusion of multiple organs. Perinatal asphyxia is an injury suffered by the fetus or NB due to poor oxygenation (hypoxia) and/or poor perfusion (ischemia) of multiple organs. Lactic acidosis is associated with hypoventilation and hypercapnia in the baby<sup>12</sup>.

Perinatal asphyxia has been defined as a delay in spontaneous breathing; low APGAR index (*Appearance Pulse Grimace Activity Respiration Effort*); cord blood acidosis; cardiotocographic registry abnormalities and also as clinical expression of post-asphyxia brain injury hypoxic-ischemic encephalopathy. There are evident signs that the prognosis after intrapartum asphyxia has improved in recent years cited by Carlos & Ermelinda, 2022<sup>13</sup>. The remaining cases, postpartum, are secondary to pulmonary, cardiovascular or neurological diseases of newborns (NB). Pathophysiological conditions that cause asphyxia include lack of oxygen from cells, excessive retention of carbon dioxide, and metabolic acidosis. Babies who have suffered the most from the asphyxia process are in a secondary apnea stage. Secondary apnea can quick-

ly cause death if the baby is not really assisted by artificial respiration and, when necessary, by compression of the heart. During secondary apnea, decreased cardiac frequency and blood pressure cause a change in skin color, from blue to white, in an effort to maximize blood flow to organs such as the heart, kidneys and glands that affect newborn children WHO, 2018, cited by Carlos & Ermelinda, 2022<sup>14</sup>.

Table 1 - Neonatal asphyxia actors (Coxim, 2015)<sup>15</sup>



amniotic fluid and pH of fetal blood. The conditions of the NB are evaluated in sequester, through three signs: respiration, heart rate c and color. The Apgar Index (*Appearance Pulse Grimace Activity Respiration Effort*) serves as a numerical and very practical report to describe the birth condition and recovery of newborns when revived by health professionals cited by Carlos & Ermelinda, 2022<sup>16</sup>.

Table. 2 - APGAR index.

Sign	0	1	2
Heart Rate	Absent	<100	>100
Respiration	Absent	Irregular	Crying hard
Muscle Tone	Stab wounds	Some bending of exterminators	Good drive
Reflexes (Nasal Stimulus)	Absent	Some movement	Sneezing
Color	Cia noise and/or pale	Rosy body and cyanotic extremities	Rosy

Apgar index value scores: Done at the 1st and 5th minute after birth<sup>17</sup>.

- 7 to 10 = NB is fine (indicates that the child has no difficulty);
- 4 to 6 = NB requires vigilance, perhaps resuscitation (indicates moderate difficulty);
- < 4 = immediate resuscitation of the NB (indicates severe suffering).

Neonatal asphyxia can be prevented with the correct interventions, in the care of the NB, according to the following steps:

Step 1: P position the baby's head correctly: Correct position can open the airways: placing the baby's head to a small extent may be enough for the baby to start breathing (do not extend or flex the neck excessively); To maintain the baby's head position, you can place a small cloth that is folded under your neck or shoulders.

Step 2: L odd airways (aspiration)

- First do the aspiration of the mouth, and only after the nose; quickly, but carefully, use a 6-8F caliber catheter and insert < 5 cm into the mouth and < 3 cm into each naring; The duration of suction must not exceed > 20 seconds; You should not suck the mucus more than twice. Note: Prolonged and deep aspiration can cause apnea.

Step 3: Ventilation

- With baby lying in the side position and with the head in extension, put on the mask,

adapted to the pumping bag, covering the baby's face: use your thumb and second finger to adjust the mask gently against the face. Use the other fingers to adjust the lower jaw of the infant gently against the mask; If the mask is not well adapted, the air comes out of the side areas of the mask. Insufflation should be carried out slowly in the proportion of 2/3 insufflations, verifying that the chest wall expands each time the pumping bag is inflated;

- Always look at the baby's chest wall during ventilation to check if the baby has made the pulmonary expansion movement through the ventilation or if it is already normally breathing alone<sup>18</sup>.


## METHODOLOGY

It uses descriptive and analytical method of quantitative approach in the application of research. Population is a collection of individual units, which can be people with one or more characteristics in common, which pertains analyze. In this research or scan that, the population corresponds to all mothers and newborns with risk factors associated with neonatal asphyxia hospitalized in the National Hospital Guido Dili Timor-Leste wingers. The sample number is 100 records whose births occurred from 1 september to 30 november 2021. The data collection instrument used for the Gilgeous observation of the midwife and the doctor. The observation grid has 5 blocks, as follows: Block I: data of

newborns, with 8 registration parameters; Block II: maternal factors, with 20 registration parameters; Block III: fetal factors, with 12 registration parameters; Block IV: labor and delivery factors, with 11 registration parameters; Hollow Belva: neonatal factors, with 9 record parameters. Data analysis we will investigate or use simple descriptive statistics to the computer program SPSS (*Statistical package For the Social Sciences*)<sup>19</sup> and the results presented in the table.

### ADOS RESULT

Graph 1. Distribution of re-born according to gender




Verification that the majority of newborns are female 53%, while 47% of newborns are male.

Graph 2 Distribution of newborns according to the diagnosis of neonatal asphyxia.




It is thought that the majority of newborns 56% presented the diagnosis of neonatal asphyxia. 25% of the sample did not present asphyxia, and 19% of the sample died before the 1st minute, according to the APGAR index.

Graph. 3 Distribution of women according to the birth of newborn.



Verification that the majority of women 61% had more than four children, with the remaining 39% four or fewer children.

Graph 4. Distribution of women according to maternal pathologies.



The most frequent and presented pathology by women was Hypertension, in 37%, followed by Pre-eclampsia/eclampsia 25%. Women with Anemia also represent 25% of the sample, and the remaining women had Diabetes 13%, based on the result investigation (2022).

Table 3. Distribution of women according to the degree of hypertension.

Hypertension	F	%
Lightweight	23	6th 2
Serious	1st	3rd 8 <sup>th</sup>
Total	37	100

It is considered that the most representative of the sample is mild hypertension, with 62% of the cases, presented hypertension grave 14 women with this diagnosis (38%).

Table 4. Distribution of women according to pre-eclampsia/Eclampsia Pre-eclampsia/eclampsia degree.

Pre-eclampsia/eclampsia	F	%
Serious	1st 15th	60
Lightweight	1st 0	40
Total	25	100

The maternal pathology pre-eclampsia/eclampsia, the one classified as Grave is the representative plus 15 (60%). However, pre-eclampsia/eclampsia Leve has a high value of 10 (40%), according the result investigation, 2022.

Table 5. Distribution of women according to the degree of anemia

Anemia	F	%
Lightweight (7 to 9g/dl)	2nd	88
Grave (< 7g/dl)	3	1st 2
Total	2nd 5th	100


We think or according that in women with anemia, anemia Leve/moderate is more frequent 22 (88%), with consequence/ grave 3 (12%).

Table 6. Distribution of women according to the classification of diabetes according to type of the cases.

Diabetes	F	%
Type I Diabetes	7	54
Gestational Diabetes	6	46
Total	13	100

I'm a record five. The time of premature rupture of membranes. Type one of diabetes 54% and gestational diabetes 46%, based on result investigation, 2022.

Graph 6. The time of premature rupture in membranes.



We felt that premature rupture of membranes more than 12 hours arose in 35 records, 51% of the sample, followed by rupture greater than 24 hours and less than 6 hours, with 25% each.

Table 7. Distribution of newborns, according to fetal factors.

Fetal factors	F	%
<b>Polyhydramnios</b>		
Yes	30	30
No	70	70
<b>Meconium</b>		
Yes	13	13
No	87	87
<b>Infection</b>		
Yes	31	31
No	69	69
<b>Newborn weight</b>		
< 1500 grams	13	13
≥ 1500 to 2499 grams	19	19
2500 to 4000 grams	68	68
<b>Birth weight unsuitable for gestational age</b>		
Yes	28	28
No	72	72

The most representative factor is 31% infection, followed by polyhydramnios with 30% of the following. Fetal factor related to Amniotic Fluid nionium appeared in 13% of the records. Regarding the weight of the NB related to gestational age, 28% presented inadequate weight, and of these 13% tin ham weight less than 1500 grams.

Table 8. The type of delivery and factors related to the umbilical cord.

Labor and childbirth	F	%
Type of childbirth		
Forceps	13	13
Cesarean	8	8
Spontaneous	78	78
Prolapse cord		
Yes	9	9
No	91	91
Short umbilical cord		
Yes	8	8
No	92	92

We found that the majority of newborns were born by spontaneous delivery 78%, followed by delivery with Forceps, 13%, and by cesarean section with 8%. In 9% of the deliveries, Prolapse of the Cord occurred, and this situation is an emergency obstetrical that requires delivery by cesarean section, and in 8% of the deliveries there was a Short Umbilical Cord, a risk factor

for the praetorium detachment of the placenta, on the basis of the research result (Carlos & Ermelinda, 2022).

Table 9. Distribution of according to Indices then APGAR at the 1st and 5th minute.

APGAR Index	Apgar indices 1 minute		Apgar indices 5 minutes	
	F	%	F	%
Lightweight (4 to 6)	2		40	59
Grave (<4)	9	38	28	41
Total	7	100	68	100

Regarding the APGAR Index at the first minute, the majority of the NB (62%), had a classification of Grave (AI<4) and 38% presented a Mild classification. At the fifth minute the APGAR Index reverted, in these two situations, to 40 records with classification moderate or Leve (59%) and 28 records (41%) with severe classification.

Table 10. Distribution of newborns, according to indices APGAR at 10 minutes.

Apgar indices 10 minutes	F	%
Record 2 to 4	3	4
Take 4 to 6	6	7
Normal 7 to 9	72	89
Total	81	100



## DISCUSSION

The moment of birth is crucial for the baby, the brain is the most delicate and noble of the 3rd response of the human body. Since this is not the case, the neonatologist has to act quickly, because the lack of oxygenation can lead to serious sequelae such as cerebral palsy or other neurologic problems, if not quickly treated. This lack of oxygenation is defined as prenatal anoxia or neonatal asphyxia adapted by Carlos & Ermelinda, 2022<sup>20</sup>. One newborn care with asphyxia may present convulsions, weight loss accentuated by difficulty in starting and maintaining breastfeeding and diffuse respiratory service. Respiratory difficulty manifests itself through a respiratory rate of less than 30 breaths per minute. Thus, it seems fundamental or basic to pay attention, especially to health personnel (both midwives and physicians), the importance of women's health surveillance, especially during pregnancy, in order to reduce the presence of pathology that may constitute a risk factor for neonatal asphyxia, such as hypertension and pre-eclampsia, providing health education to mothers to avoid neonatal asphyxia cited by Carlos & Ermelinda, 2022<sup>21</sup>.

Hypertension is the most common cardiovascular disease during pregnancy and even during the fertile years of women <sup>22</sup>. Complications resulting from hypertensive disease are, alongside infection and bleeding, the main cause of maternal death in most specialized services. Gestational pretensions correspond to cases of increased blood pressure, without proteinuria, after the 20th week of pregnancy, returning the blood levels elevated to normal in the postpartum period (12 weeks)<sup>23</sup>. Late or transient events or gestational hypertension, defined as the development of increased pressure values, occurring antepartum, labor or in the first 24 hours postpartum, without other signs of preeclampsia or pre-existing hypertension. Normal blood pressure should return to normal within 10 days after delivery<sup>24</sup>.

Among all the pathologies that manifest or worsen during pregnancy, preeclampsia/eclampsia is the most frequent and the one that is accompanied by higher maternal and perinatal morbidity and mortality, characterized by the presence of hypertension, proteinuria and edemas, in addition to other alterations. In developing countries such as East Timor, hypertension in pregnancy, broad or broad sense, is one of the main causes of mortality. According to the data obtained in this research, severe preeclampsia/eclampsia presented 60% of the total of the 25 cases. Mildrécampsia, diastolic blood pressure is less than 100 mmHg, proteinuria is 1+ or 2+ in the reagent strip, many women have edema in the face and hands and weight gain is observed. For Sousa (2017) severe preeclampsia presents several signs and symptoms that indicate the severity of

preeclampsia, there are criteria that define it as 90% of the total iron requirement is used in the those proposed by Picher and MacDonald last trimester of pregnancy. Counseling, education and help is needed for pregnant women to understand this need. Anemia in pregnant women can be defined as a condition in which hemoglobin levels are below normal 11g %, which can cause hypoxia and reduced blood flow to the uterus, which will lead to reduced oxygen flow to the placenta and fetus, and can use interruption of breathing. Diabetic patients are at high-risk hypoglycemia the first three days of the newborn, even when they eat well. Our sample presented 7 participants with type 1 diabetes (54%), and 6 with gestational diabetes (46%). To reduce the impact on the mother, before conception, a careful evaluation of the presence of chronic complications of diabetes should be carried out, as some may lead to increased morbidity for the pregnant woman. On the other hand, the abnormal metabolic environment caused by hyperglycemia has a significant impact on pregnancy and the fetus or baby to have carefully, so this whole situation should be treated and controlled before pregnancy<sup>26</sup>.

When the newborn baby (NB) has mild breathing difficulties, interventions or interferences to prevent neonatal asphyxia include maintaining heat, positioning the baby's head correctly, cleaning the breathing pathways (aspiration) and allowing amniotic and breastfeeding or breastfeeding to be important. If the difficulty is maintained and suffocation is severe it may be necessary to provide assisted ventilation. These inter-

As the situation increases, cyanosis, pulmonary edema, pain in the right hypochondria, thrombocytopenia (<100,000), microangiopathic hemolytic anemia, jaundice or abnormal liver function may occur, and the pregnant woman may suffer from seizures. Specific competencies to prevent and treat this pathology include; identifying the risk factors for preeclampsia and eclampsia, obstetric and fetal development, and tough care for seizure (Carlos & Ermelinda, 2022). In relation to prenatal consultations, there was a greater prevalence of NEWBORNS with asphyxia among mothers who underwent up to 4 pre-natal 2<sup>5</sup> Preeclampsia and eclampsia as a complication of hypertension that is present before pregnancy, or diagnosed before 20 weeks of gestation, which makes it imperative to control hypertension throughout pregnancy and before childbirth 2<sup>5</sup>. In the sample studied verify, 88% of mild anemia was found, and this is a known pathology, not those pregnant in East Timor. In pregnancy with a single fetus, maternal needs range from 800 to 1,000 mg of iron, from 300 to 350 mg for the formation of the placental fetus unit, in addition to the amount available to expand the maternal hemoglobin mass. About

ventions should be validated with the pediatrician or neonatologist, who should perform a complete examination of the NB to establish the therapeutic plan that best appropriates the situation<sup>27</sup>. In the present study, the records reveal that at the first minute, the majority of newborns (62%) had a Severe classification with APGAR index below 4, and 38% presented a Mild classification (score between 4 and 6). However, in the five-minute evaluation, this situation reversed, moving to 40 records with a Level rating (59%) and 28 records (41%) with a Severe rating. That is, after adequate intervention to support the vital functions of the NB, particularly with regard to the respiratory, there was a recovery of the APGAR score. When we analyzed the records of the APGAR index evaluation at 10 minutes, we found that most of the NB (89%) already had a normal score of 7/9, according to the scanning result (2022).

## VI CONCLUSION

Perinatal asphyxia is an injury suffered by the fetus or newborn (NB), due to poor oxygenation or poor perfusion of multiple organs. From the results of this study, comparing with the references in the literature, the factors associated with the occurrence of neonatal asphyxia were identified in this group of newborns: multiparity, hypertension, preeclampsia/eclampsia, anemia, premature rupture of membranes greater than 12 hours, polyhydramnios and in-

The diagnosis of neonatal asphyxia is made from the evaluation or evaluation of the five parameters that constitute the APGAR Index (Heart Rate; Breathing; Muscle Tone; Reflexes; Color), scored from 0 to 2 according to the characteristics of each parameter that the newborn presents. As mentioned above, a 9/10 APGAR index score at the first minute is considered optimal and reveals an NB without respiratory, cardiac and neuromuscular difficulties. Considering the pattern referenced, the data obtained in our study reveal that, unfortunately, there is still a high number of newborns who have an APGAR index between 4 and 6, at the 1st minute of life, which places them in the classification of severe to mild asphyxia. Although we recognize that, at the 5th minute of life there was a recovery in the APGAR score (reduction of 34 records in the Severe classification and increase of two records in the Level classification) and that at the 10th minute most newborns already had a score of 7/9, we have to mourn the death of 19% of the newborns, who died before the 1st minute according to the records obtained, probably related to complications of pregnancy and/or childbirth, which makes it urgent to identify and establish concrete strategies aimed at reducing neonatal and infant mortality, there is national and international level according to this study (Carlos & Ermelinda, 2022).

## REFERENCES

1. WHO, (2019). *Conduct in Newborn Problems. Nurses and Midwives, Department of Reproductive Health and Research*. Geneva: a Guide to (Doctors. Monica Translation of Queiroz Tels Saponi Neves),- ISBN-13:978-85-7241-618-4:São Paulo: ROCA. <http://www.who.int>
2. Moreira, (2015). *The high-risk newborn: theory and practice of care*. Rio de Janeiro: Fiocruz.
3. Orhan (2010). *Nursing in the health of women, mothers and newborns: lifelong care*. Porto Alegre.
4. WHO (2015). *Eclampsia-Education for a Safe Maternity: modules of education*. Brazil, 6nd edition.
5. Joseph, A. (2012). *Statistics of CPLP*, INE, IP. Lisbon, Portugal.
6. Ministry of Health (2019). *Book Assistance Safe Childbirth*, East Timor.
7. Ministry of Health. (2018). *Health Statistics Report*, Health Information System Office, East Timor.
8. Ministry of Health. (2016 to 2017). *Health Statistics Report*, Health Information System Office, East Timor.
9. Guido Valadares National Hospital. (2019). *Maternal and Child Health Statistics Report*. Dili East Timor.
10. UNICEF, (2011). *Safe Childbirth Care Learning Book*. Ministry of Health, East Timor.
11. Gonzalez, (2010). *Perinatal asphyxiation*. Pediatric Nursing Practice.
12. Zanonía, (2014). *Perinatal Asphyxia Chapter of the book Assistance to the Newborn at Risk*, edited by Paulo R. Marlite: 2nd and 4th Edition
13. Sandra and Pires. (2012)., *Perinatal Asphyxia and Maternal Behavior*, Psychological Analysis., 3 (XX): 439-448.
14. Souza, M.F. (2013). *Factors associated with perinatal asphyxia: a population study based on the Live Birth Information System*. Rio de Janeiro, RJ: IFF/ FOC. Brazil.
15. Cushion, (2015). *Asphyxia intervention in Newborn for Midwife*. The Documents of the Ministry of Health.
16. Marotta and Rezende, (2017), *Prevalence of perinatal asphyxia*. Rio de Janeiro. Cad. Public Health, v. 17, 2015.
17. Lowdermilk Perry, (2016). *Maternity nursing*. Sedate Edition, copyright @ 2016 2013, by Mosby Inc. Editor: Catherine Jackson. ISBN-13:978-323-03366-4.
18. UNICEF, (2015). *Book Assistance in Neonatal Nursing.Adaptation in Timor-Leste: Module I, II, III and IV*, Ministry of Health, East Timor,
19. Tilman C.B et al. (2020). *The Perception of Population and Health Professionals regarding the National immunization Program of Timor-Leste*. Health Systems and Policy Research, ISSN 2254-9137 Vol.7 No.1:2 2020.

- [www.imedpub.com](http://www.imedpub.com) published date May 11, 2020.
20. Lath E., et al (2011), *Apgar test scores difference between children born of normal delivery and cesarean delivery*. Art. Unipara Health Sciences, 5(3): 211
21. Canal, A. (2014). *Factors associated with perinatal asphyxia*. Rio Janeiro (Rio J), RBCO v.26, n.10.
22. Baraco, A. (2014). *Prevention of evidence-based preeclampsia*. Rio de Janeiro, 6<sup>2</sup> (7) Id: 521744.
23. Montenegro E, (2016). *Gestational preten-sion*. Guide to Effective Care in Pregnancy and Childbirth. Rio de Janeiro, 2015, 3rd edition, Guanabara Kagan.
24. Faraz ani S. (2010). *Hypertension. Pregnant women*. Saudis J Kinder Transpolar 644<sup>3</sup>; 10:298-312.
25. Sousa, P, (2017). *Study of maternal risk factors associated with neonatal sepsis early in Brazilian Amazon Hospital*. Online version ISSN 1806-9339. Rev. Gingerol Bras. Octeto. Vol.29.no.8 Rio de Janeiro Aug.2007. <http://dx.doi.org/10.1590/S0100-7203200700080002>.
26. Nascimento L. e Ramos, (2014). *Application of the CRIB score as a predictor of death in a neonatal intensive care unit: an expanded approach*. Mater Health. Infant, 4(2):151-7.
27. Pinhel, S. (2018), *Prevalence of Perinatal Asfixia*, Hospital Municipal São Lucas, 645<sup>1</sup>. Revista Saúde Coletiva, n.2.

**Citation:** Carlos Boavida Tilman, et al. FACTORS ASSOCIATED WITH THE INCIDENCE OF NEONATAL ASPHYXIA IN NEWBORNS AT THE GUIDO VALADARES NATIONAL HOSPITAL DILI TIMOR-LESTE AJMCRR. 2022; 1 (2): 1-13.